

## The Abbey Pain Scale

*For assessment of pain in patients who cannot verbalise i.e. patients with dementia or communication difficulties*

### **Use of the Abbey Pain Scale**

The Abbey Pain Scale is best used as part of an overall pain management plan.

### **Objective**

The Pain Scale is an instrument designed to assist in the assessment of pain in patients who are unable to clearly articulate their needs.

### **Ongoing assessment**

The Scale does not differentiate between distress and pain, so measuring the effectiveness of pain-relieving interventions is essential.

Recent work by the Australian Pain Society recommends that the Abbey Pain Scale be used as a movement-based assessment.

The staff recording the scale should therefore observe the patient while they are being moved, e.g. during pressure area care, while showering etc.

Complete the scale immediately following the procedure and record the results on the Abbey Pain tool chart.

Include the time of completion of the scale, the score, staff member's signature and action (if any) taken in response to results of the assessment, e.g. pain medication or other therapies.

A second evaluation should be conducted one hour after any intervention taken in response to the first assessment, to determine the effectiveness of any pain-relieving intervention.

If, at this assessment, the score on the pain scale is the same, or worse, consider further intervention and act as appropriate.

Complete the pain scale hourly, until the patient appears comfortable, then four-hourly for 24 hours, treating pain if it recurs.

Record all the pain-relieving interventions undertaken. If pain/distress persists, undertake a comprehensive assessment of all facets of patient's care and monitor closely over a 24-hour period, including any further interventions undertaken.

If there is no improvement during that time, notify the medical practitioner of the pain scores and the action/s taken.

## ABBEY PAIN ASSESSMENT SCALE (follow on assessment)

	Date and time									
<b>VOCALISATION</b>										
e.g.: whimpering, groaning, crying <i>Absent 0 Mild 1 Moderate 2 Severe 3</i>										
<b>FACIAL EXPRESSION</b>										
e.g.: looking tense, frowning grimacing, looking frightened <i>Absent 0 Mild 1 Moderate 2 Severe 3</i>										
<b>CHANGE IN BODY LANGUAGE</b>										
e.g.: fidgeting, rocking, guarding part of body, withdrawn <i>Absent 0 Mild 1 Moderate 2 Severe 3</i>										
<b>BEHAVIOURAL CHANGE</b>										
e.g.: increased confusion, refusing to eat, alteration in usual patterns <i>Absent 0 Mild 1 Moderate 2 Severe 3</i>										
<b>PHYSIOLOGICAL CHANGES FACIAL</b>										
e.g.: temperature, pulse or blood pressure outside normal limits, perspiring, flushing or pallor <i>Absent 0 Mild 1 Moderate 2 Severe 3</i>										
<b>PHYSICAL CHANGES</b>										
e.g.: skin tears, pressure areas, arthritis, contractures, previous injuries <i>Absent 0 Mild 1 Moderate 2 Severe 3</i>										
TOTAL score =										
Signature of person completing score:										
0-2 No pain	3-7 Mild pain			8-13 Moderate Pain			14+ Severe Pain			

The Pain Scale is an instrument designed to assist in the assessment of pain in patients who are unable to clearly articulate their needs.

The Abbey pain scale does not differentiate between distress and pain, therefore measuring the effectiveness of pain-relieving interventions is essential.

The pain scale should be used as a **movement-based assessment**, therefore observe the patient while they are being moved, during pressure area care, while showering etc.

A second evaluation should be conducted **1 hour after** any intervention taken. If, at this assessment, the score on the pain scale is the same, or worse, consider further intervention and act as appropriate.

Complete the scale hourly until the patient scores mild pain then 4 hourly for 24 hours treating pain if it recurs.

If the pain/distress persists, undertake a comprehensive assessment of all facets of the patients care and monitor closely over 24 hours including further intervention undertaken.

If there is no improvement during that time notify the doctor/pain team of the pain scores and actions taken.

# Abbey Pain Scale

For measurement of pain in patients who cannot verbalise

Person completing the scale:

**Name:** ..... **Designation:** .....

*How to use scale: While observing the patient score questions 1-6*

**Q1. Vocalisation**

e.g., whimpering, groaning, crying

Absent 0 Mild 1 Moderate 2 Severe 3

**Q1**

**Q2. Facial expression**

e.g., looking tense, frowning grimacing, looking frightened

Absent 0 Mild 1 Moderate 2 Severe 3

**Q2**

**Q3. Change in body language**

e.g., fidgeting, rocking, guarding part of body, withdrawn

Absent 0 Mild 1 Moderate 2 Severe 3

**Q3**

**Q4. Behavioural change**

e.g., increased confusion, refusing to eat, alteration in usual patterns

Absent 0 Mild 1 Moderate 2 Severe 3

**Q4**

**Q5. Physiological change**

e.g., temperature, pulse or blood pressure outside normal limits, perspiring, flushing or pallor

Absent 0 Mild 1 Moderate 2 Severe 3

**Q5**

**Q6. Physical changes**

e.g., skin tears, pressure areas, arthritis, contractures, previous injuries

Absent 0 Mild 1 Moderate 2 Severe 3

**Q6**

**Add Scores 1 – 6 and record here**

Total pain score =

**Now tick the box that matches the total pain score**

0 – 2

3 – 7

8 – 13

14 +

No pain

Mild

Moderate

Severe

**Finally, tick the box which matches the type of pain**

Acute

Chronic

Acute  
&  
Chronic