



CLINICAL GUIDELINE

Pulmonary Embolism, Investigation and Management

A guideline is intended to assist healthcare professionals in the choice of disease-specific treatments.

Clinical judgement should be exercised on the applicability of any guideline, influenced by individual patient characteristics. Clinicians should be mindful of the potential for harmful polypharmacy and increased susceptibility to adverse drug reactions in patients with multiple morbidities or frailty.

If, after discussion with the patient or carer, there are good reasons for not following a guideline, it is good practice to record these and communicate them to others involved in the care of the patient.

Version Number:	4
Does this version include changes to clinical advice:	Yes
Date Approved:	4 th May 2023
Date of Next Review:	31 st May 2026
Lead Author:	Colin Church
Approval Group:	Medicines Utilisation Subcommittee of ADTC

Important Note:

The Intranet version of this document is the only version that is maintained. Any printed copies should therefore be viewed as 'Uncontrolled' and as such, may not necessarily contain the latest updates and amendments.

Investigation and Management of Pulmonary Embolism

Date March 2023

Authors Dr Colin Church (Respiratory and Pulmonary Vascular Consultant, QEUH)
Dr Harrison Stubbs (Pulmonary Vascular Fellow, GJNH)

Contents

Remit.....	1
Quick Definitions.....	1
Introduction	2
Pre-Diagnostic Scores and Diagnosis	3
Severity Assessment	5
Management of High-Risk PE (Massive PE)	6
Management of Intermediate-risk PE.....	10
Management of Low-Risk PE	11
Anticoagulation.....	13
Duration of Anticoagulation	16
Discharge and Follow Up	17

Remit

This guideline is designed for use in non-pregnant adults in Secondary Care. It has been designed to be a comprehensive overview for the diagnosis and treatment of pulmonary embolism in this population. Separate guidelines are available for use in pregnant patients.

Quick Definitions


	High Risk PE	Haemodynamic instability - sustained hypotension (SBP <90mmHg, for 15 minutes) or requiring inotropic support, that is not related to a cause other than PE (i.e. arrhythmia, LV dysfunction, sepsis or hypovolaemia).
	Intermediate-High Risk PE	Haemodynamically stable - but at risk of rapidly deteriorating with evidence of right ventricular dysfunction (on CTPA or TTE) <u>and</u> myocardial injury (i.e. raised troponin).
	Intermediate-Low Risk PE	Haemodynamically stable – lower risk than intermediate high-risk but clinical markers of PE severity and RV dysfunction may still be present.
	Low Risk PE	Haemodynamically stable – no evidence of clinical markers of severity, no RV dysfunction and no myocardial injury.

Table 1. Quick Definitions for Risk Stratification in Pulmonary Embolism

Introduction

Pulmonary embolism (PE) occurs when emboli, arising from a blood clot in the venous system, obstruct the pulmonary arterial system, leading to respiratory and cardiovascular dysfunction.

- PE should be systematically stratified and managed according to the risk of haemodynamic compromise.
- Patients with suspected PE with haemodynamic compromise should be treated urgently, as for High Risk PE.
- In patients with suspected PE who are haemodynamically stable, pre-diagnostic predictive scores should be used to assess the need for further investigations.
- Patients with confirmed PE should undergo severity assessment to determine the management and bed allocation.
- All patients with PE should have a follow up in a PE clinic.
- The following algorithm outlines the general approach to a patient with suspected PE. For details see relevant sections below.



*Haemodynamic instability - sustained hypotension (SBP <90mmHg, for 15 minutes) or requiring inotropic support, that is not related to a cause other than PE (e.g. arrhythmia, LV dysfunction, sepsis or hypovolaemia)

Figure 1. Overview of investigation and treatment of pulmonary embolism

Pre-Diagnostic Scores and Diagnosis

- If a patient has a suspected PE with haemodynamic instability (SBP<90) refer to **High Risk PE**.
- Patients who are stable should have PE considered based on predictive scores and consideration for further testing – see Figure 2.

Signs and Symptoms

- Chest pain (often pleuritic), cyanosis, dyspnoea, haemoptysis, syncope
- Tachycardia, hypotension, raised JVP, hypoxaemia, tachypnoea

Assess Predictive Score

1. If the patient is haemodynamically unstable go straight to **High Risk PE**.
2. If in ED or medical receiving: for low risk patients the Pulmonary Embolism Rule Out Score (PERC) can be used to rule out PE.
3. Perform a pre-predictive calculation (such as the revised Geneva score or the Wells score) to assess the probability of PE and inform investigations.

PERC Score (Pulmonary Embolism Rule Out Score)

This is used for patients with a low likelihood of PE, who demonstrate low-risk features, and can be used to rule out PE in the Assessment Unit or the Emergency Department.

A score of ≥ 1 means PE **cannot** be ruled out.

Age ≥ 50	1
Heart Rate ≥ 100	1
Peripheral oxygen saturation $\leq 95\%$	1
Unilateral leg swelling	1
Haemoptysis	1
Surgery or trauma ≤ 4 weeks ago requiring general anaesthetic	1
Previous PE or DVT	1
Hormone Use e.g. oral contraceptive, hormone replacement or oestrogen containing hormones	1

Table 2. Pulmonary Embolism Rule Out (PERC) Score

Revised Geneva Score

Assesses the probability of PE in order to inform investigations.

Age ≥ 65 years	1
Previous DVT/PE	3
Recent surgery or lower limb fracture (≤ 1 month)	2
Malignant disease (active or cured ≤ 1 year)	2
Unilateral lower limb pain	3
Haemoptysis	2
Heart rate 74 – 94 bpm	3
Heart rate ≥ 95 bpm	5
Pain on deep venous palpitation of leg and unilateral oedema	4

Table 3. Revised Geneva Score

Score	Probability of PE	Incidence of PE
0-3	Low	~7-9%
4-10	Moderate	~20-30%
≥ 11	High	>60%




Figure 2. The use of scoring systems to predict pulmonary embolism in patients without haemodynamic instability.

Severity Assessment

All confirmed PE should be stratified and managed according to the severity. This is judged by;

1. Haemodynamic instability (shock) - hypotension
2. Myocardial damage – troponin biomarkers
3. Right ventricular strain – RV size and function on CTPA or TTE
4. Risk of early mortality – Simplified PE Severity Index (sPESI)




Figure 3. Severity assessment in pulmonary embolism

sPESI Score Simplified Pulmonary Embolism Severity Index

Age ≥ 80	1
History of cancer	1
History of cardiorespiratory disease	1
Heart rate ≥110 bpm	1
Systolic BP ≤100 mmHg	1
Peripheral O2 saturations ≤90%	1

Table 4. Simplified Pulmonary Embolism Severity Index (sPESI)

Management of High-Risk PE (Massive PE)

1. High-risk PE is defined as an acute PE with sustained hypotension (SBP <90mmHg, for 15 minutes) or requiring inotropic support, that is not related to a cause other than PE (e.g. arrhythmia, LV dysfunction, sepsis or hypovolaemia).
2. Death occurs as increased right ventricle afterload leads to strain placed on the right ventricle, causing a reduction in cardiac output, hypotension and hypoxaemia.

Investigations

1. Confirmation by CTPA is preferable and also allows assessment of pulmonary vascular anatomy and RV assessment
2. If not possible due to clinical condition (e.g. intubated) urgent bedside TTE should be performed to assess for RV size and dysfunction

Management

- Move to level 2 or level 3 area (resus, HDU, CCU, ITU) for monitoring and treatment
- Give resuscitation care which may include appropriate oxygen therapy, invasive blood pressure monitoring, fluid resuscitation and inotropic support.
- If PE confirmed, or felt highly likely, give **reperfusion therapy**, which in most centres will currently consist of systemic thrombolysis.
- Full-dose Systemic thrombolysis should be the first consideration, but if there are concerns of a high bleeding risk, or systemic thrombolysis has been unsuccessful, other modalities should be considered.

Summary of Reperfusion Therapy

Reperfusion Therapy			
Systemic Thrombolysis (Full Dose)	Systemic Thrombolysis (Low Dose)	Catheter-directed Thrombolysis (CDT)	Surgical Embolectomy
Considered as first line for treatment of High Risk PE	Considered in High Risk PE or Intermediate-High Risk PE – where it is felt the bleeding risk outweighs the benefit of full dose systemic thrombolysis	Considered when systemic thrombolysis has been unsuccessful or in High Risk and Intermediate-High Risk PE – where it is felt the bleeding risk outweighs the benefit of either full-dose or low-dose systemic thrombolysis	Considered when; <ul style="list-style-type: none"> - Systemic thrombolysis has been unsuccessful - High Risk and Intermediate-High Risk PE – where it is felt the bleeding risk outweighs the benefit of either full-dose or low-dose systemic thrombolysis - Transient clot in the RA, RV or traversing a PFO to reduce the risk of rapid deterioration or stroke (in the case of a PFO).

Table 5. Overview of options for reperfusion therapy in pulmonary embolism

Systemic Thrombolysis (Full Dose)

- Systemic thrombolysis **is** safe to give if LWMH has already been administered.
- Thrombolysis is more effective within the first 48 hours of presentation, yet can be considered at any time especially if there is subsequent clinical deterioration.
- There is no evidence to suggest that thrombolysis decreases the subsequent risk of Chronic Thromboembolic pulmonary hypertension (CTEPH). A high clot burden is not an indication, in itself, for thrombolysis.
- Consider contra-indications to thrombolysis;

Absolute*	Relative*
Haemorrhagic stroke (at any time)	TIA in last 6 months
Ischaemic stroke in last 6 months	Pregnancy or within 1-week post-partum
CNS damage or neoplasia	Advanced liver disease
Major trauma/surgery/head injury in last 3 weeks	Infective endocarditis
GI Bleeding within the last month	Active peptic ulcer

Table 6. Contra-indications to thrombolysis

*In High-Risk PE, even in the presence of contra-indications, it may be that thrombolysis is still considered and given, due to the high risk of death. However this is a decision for senior clinicians and should be first discussed with the responsible consultant.

If possible, **consent** that patient for thrombolysis. Relative risks of thrombolysis are shown below.

	Thrombolysis + Anticoagulation	Anticoagulation alone
Major Bleeding	~10-15%	~4%
Intracranial Haemorrhage	~2%	~0.2%
Death from all causes	~2%	~4%

Table 7. Relative risks of thrombolysis and/or anticoagulation in High-Risk PE

Whilst these can be significant risks, they need to be taken in the context of patient's clinical condition. Systemic thrombolysis is strongly recommended if the criteria for High-Risk PE are met.

Alteplase

- The IV Monograph for Alteplase can be found on the Medusa website.

Alteplase Dose	Patients ≥ 65kg	Patients < 65kg	Peri-arrest or cardiac arrest
IV Bolus (over 2 minutes)	10mg	10mg	50mg
Immediately followed by an IV Infusion (over 2 hours)	90mg	Maximum dose 1.5mg/kg	50mg
Note	In total 2x50mg vials of Alteplase	e.g if 50kg, infuse 75mg over 2 hours after bolus	

Table 8. Dose of Alteplase in full dose systemic thrombolysis

- If alteplase is not available, consider other thrombolytics such as tenecteplase or follow local MI thrombolysis regimen (note these are unlicensed for use in PE).

Unfractionated Heparin (UFH)

- Upon completion with alteplase, commence UFH. Only use 1000units/ml concentration. If there are concerns with UFH, subcutaneous LMWH may be considered.
- If the patient has had treatment with LMWH within 12 hours, do not give a bolus dose.
- If no LMWH within 12 hours **and** the patient weighs between 50-100kg;
 - Give a **bolus of 5000 units UFH IV over 5 minutes**.
 - Followed by **maintenance infusion** of UFH at a rate of 18units/kg/hour
e.g. usually ~ 1,200units per hour for a 70kg patient.
 - If patient is at high risk of bleeding, start at 1000units/hour.
- If the patient weighs <50kg or >100kg – see the heparin dose adjustment guideline ([link](#)).
- Check APTT ratio after 6 hours – aiming 1.8-2.8. Adjust as per table below. If an adjustment is made, recheck APTT ratio after 4 hours. If no change, check APTT ratio daily.

UFH Dose Adjustment	
APTT Ratio	UFH Infusion Rate Change
>4	Stop for 60 minutes and recheck APTT ratio, before recommencing at a rate reduced by 300–500units/hour
3.5–4	Stop for 60 minutes and reduce heparin by 200units/hour
2.9–3.4	Stop for 30 minutes and reduce heparin by 100units/hour
1.8–2.8	TARGET - No change
1.2–1.7	Increase heparin by 200units/hour
<1.2	Increase heparin by 400units/hour and consider further bolus of 5,000units heparin
- Check APTT ratio every 4 hours after any adjustment in infusion rate. - Patients rarely require rates >1.6-2ml/hour. If target APTT ratio is not achieved at this rate, monitor anti-Xa levels (target 0.35-0.7units/ml). - Routine platelet monitoring for HIT is not required unless <3 months from recent surgery.	

Table 9. Dose adjustments of unfractionated heparin based on APTT Ratio

Further Management

UFH can be stopped once the patient is deemed to be stable. Ongoing anticoagulation should be commenced. The choice of ongoing anticoagulation is discussed in detail below.

Systemic Thrombolysis (Low Dose)

Low Dose Systemic Thrombolysis should be considered when;

1. High Risk PE or Intermediate-High Risk PE – where it is felt the bleeding risk outweighs the benefit of full dose systemic thrombolysis

It is administered by;

1. **Alteplase 10mg IV bolus over 2 minutes,**
2. Followed by **40mg Alteplase IV infusion over 1 hour**
3. Commence UFH as for High Risk PE. Aim for the same APTT target of 1.8-2.8

Catheter Directed Thrombolysis (CDT)

CDT should be considered when;

1. Systemic thrombolysis has been unsuccessful
2. High Risk and Intermediate-High Risk PE – where it is felt the bleeding risk outweighs the benefit of either full-dose or low-dose systemic thrombolysis

CDT uses a lower dose of thrombolysis and has a safer bleeding risk profile than systemic thrombolysis. However, it takes time to set-up and may not be available immediately.

Direct clot retrieval is not currently available, but may become available in the future.

This technique is organised through contacting Interventional Radiology. A pulmonary artery catheter is placed into the right and left Pulmonary artery and alteplase is infused at 10mg/24 hrs into each. If CDT is felt to be the most suitable therapy, but is not available at your site, site transfer should be considered.

Surgical Embolectomy

Surgical embolectomy should be considered when;

1. Systemic thrombolysis has been unsuccessful
2. High Risk and Intermediate-High Risk PE – where it is felt the bleeding risk outweighs the benefit of either full-dose or low-dose systemic thrombolysis
3. Transient clot in the RA, RV or traversing a PFO to reduce the risk of rapid deterioration or stroke (in the case of a PFO).

Surgical embolectomy requires a careful MDT approach including ITU, anaesthetics, cardiothoracic surgery and haematology. Potential surgery may still be considered after recent thrombolysis is administered.

Mechanical Circulatory Support in High Risk PE

The use of advanced mechanical circulatory support should be assessed on an individual case basis. It should only be considered if there is the potential for further reperfusion intervention or to support a patient once reperfusion therapy has been instituted and there is clinical agreement that recovery could still occur. This would require discussion with the local ICU and Golden Jubilee ICU team.

Management of Intermediate-risk PE

Intermediate-risk PE patients may present with significant symptoms and/or large volume clot on CTPA whilst not meeting high risk criteria. These patients may have a positive troponin (indicating myocardial damage), a high risk sPESI or evidence of RV dysfunction on imaging. Intermediate-risk patients have a higher mortality than low-risk patients and can deteriorate rapidly, warranting consideration for level 2 monitoring and admission for observation.

Management

- If hypotensive (SBP <90mmHg) refer to the **High-risk PE** guideline

Intermediate-High Risk

Patients with Intermediate-High Risk PE are at risk of rapidly deteriorating, most often within 48 hours of admission. Therefore;

- Monitor in a level 2 bed (e.g. HDU, CCU or an acute care environment with cardiac monitoring and adequate nursing provision)
- Monitor as an inpatient for a minimum of 48 hours
- Treat with LMWH for a longer period (i.e. 2-3 days) before converting to DOAC
- If at presentation patients appear critically unwell (e.g. marked hypoxaemia, high lactate, signs of right heart failure) but not meeting criteria for High Risk PE, **reperfusion therapy** should still be considered.
- If clinically deteriorating (e.g. progressive hypoxaemia, rising heart rate, high or rising lactate, clinical features of circulatory compromise e.g. cold, clammy, cyanosis), consider **reperfusion therapy** in any modality.

See the details on these treatment options under High Risk PE

- Whilst all modalities should be considered and be appropriate to the patient, low-dose thrombolysis perhaps provides the lowest chance of dying from PE whilst simultaneously reducing the bleeding risk.

Intermediate Low-Risk

These patients should be monitored on a ward.

- Once PE is confirmed on CTPA, LMWH can be stopped and a DOAC commenced for acute and ongoing treatment (see Anticoagulation section).
- Patients should be monitored as an inpatient for a minimum of 24 hours, yet discharge within 48 hours of presentation should be guided by senior clinical judgement.

Management of Low-Risk PE

- Patients with low risk PE should be assessed for early discharge.
- Use the below flowchart to assess suitability for outpatient management.




Figure 4. Overview for the ambulatory management of suspected low risk pulmonary embolism.

Suitable for Early Discharge/Ambulatory Care

- Low Risk PE
- No severe chest pain
- No significant co-morbidity that may complicate acute PE and/or its treatment
e.g. significant renal/liver/cardiac dysfunction
- No other requirement for admission
- Patient is mobile and able to return to hospital
- Patient is happy for ambulatory management
- Patient is aware of the need to return to hospital earlier if they become more unwell at home (should be warned of presyncope symptoms, becoming more breathless, etc.)
- Patient is **not** pregnant (see Special Cases)
- Patient is **not** already therapeutically anticoagulated.
- Bleeding risks and contra-indications to LWMH have been ruled out
- Patient does not live alone or does not have good support at home
- Patient has a working telephone

Ambulatory Management of Low Risk PE

- Give treatment dose LMWH or DOAC
- Ensure the patient has been educated on their new anticoagulant and how to self administer.
- The GGC DOAC booklet can be given to patients and can be found here: [“GGC Medicines: Direct Oral Anticoagulation \(DOAC\) Patient Information Booklet and Alert Card”](#)
- Request CTPA
- Ensure the patient has enough doses of LMWH/ DOAC to administer at home until the CTPA.
- Explain plan to patient– ensure they understand follow-up, CTPA and LMWH/ DOAC.
- Request follow-up as per local protocol e.g. hot clinic, ambulatory care pathway.

Anticoagulation

The aim of anticoagulation is to complete the treatment of the acute PE and prevent recurrence of VTE over the long-term.

Subcutaneous Low Molecular Weight Heparin (LMWH)

Dalteparin is the LMWH of choice across NHS GGC unless the patient is pregnant or has specific contraindications to dalteparin.

Continue dalteparin until:

- The diagnosis is disproved
- The diagnosis is confirmed – switch patient to oral anticoagulation (see relevant sections for DOACs and warfarin)

Dalteparin Dose

Actual weight (kg)	Dalteparin once daily dose (units)	Pre-filled syringe colour
34-45	7,500	Green
46-56	10,000	Red
57-68	12,500	Brown
69-82	15,000	Purple
≥83	18,000 (maximum dose)	Grey

Renal Impairment and Extremes of Body Weight

Guidance on dose adjustment for patients with significant [renal impairment](#) or [weighing >120kg](#) are available on the [GGC Clinical Guidelines Platform](#) and in the [GGC Therapeutics Handbook](#).

Unfractionated Heparin (UFH)

- See guidance under High Risk PE.
- Used in treatment of DVT / PE when:
 - rapid anticoagulation is deemed appropriate (e.g. high-risk PE)
 - patients are at particularly high bleeding risk (e.g. recent surgery/trauma),
 - when anticoagulation may need to be urgently reversed.

Direct Oral Anticoagulants (DOACs)

Apixaban is the DOAC of choice across NHSGCC for acute treatment and long-term secondary prevention of PE.

Rivaroxaban, Edoxaban and Dabigatran remain on formulary for special cases when they may be preferable. All DOACs are as effective as warfarin for the treatment and prevention of PE and are associated with a lower bleeding risk yet do not require monitoring. Reversal agents for life-threatening haemorrhage are available on the [GGC Clinical Guideline Platform](#) (after discussion with haematology) for dabigatran (idarucizumab [Praxbind]), apixaban and rivaroxaban (andexanet alfa [Ondexxya]) and off-licence for edoxaban (andexanet alfa).

Apixaban Contra-indications

- Apixaban should not be used if the Creatinine Clearance is <15ml/minute and used with caution if 15-29mls/min. Further information can be found on the GGC Clinical Guideline platform ([link](#)).
- Liver disease, associated with cirrhosis and/or coagulopathy
- Pregnancy or breast feeding
- Concurrent treatment with azoles (except fluconazole), protease inhibitors, or other cytochrome P450 inducers (e.g. Rifampicin, Phenytoin, Carbamazepine, Phenobarbitol)

Treatment Regimen

- Treat with LMWH until the diagnosis is objectively confirmed.
- If the diagnosis has already been confirmed (e.g. an incidental finding on a CT scan) and low-risk, then treatment can immediately start with a DOAC.
- Start Apixaban 22-24 hours after the last dose of LMWH
- Loading dose – **Apixaban oral, 10mg twice daily, for 7 days**
- Maintenance dose – **Apixaban oral, 5mg twice daily** for duration of anticoagulation
- **If** long term anticoagulation, after 3-6 months reduce to Apixaban oral 2.5mg twice daily.

Discharging on Apixaban

- The GP should be informed on the initial duration of anticoagulation using the IDL.
- The GGC DOAC booklet can be given to patients and can be found here: "[GGC Medicines: Direct Oral Anticoagulation \(DOAC\) Patient Information Booklet and Alert Card](#)"

Renal impairment and DOACs

- Renal impairment affects the use and dosage of DOACs.
- Creatinine clearance (CrCl) must be [calculated](#) as eGFR is not reliable for DOAC dosing.
- For Creatinine Clearance <15mls/min, DOACs cannot be used and LMWH or Warfarin should be used.
- Apixaban should be used first line for all patients, if Creatinine Clearance is >15mls/min

Creatinine Clearance (ml/min)	Apixaban	Rivaroxaban	Dabigatran	Edoxaban
30-50	No dose adjustment	Reduce dose to 15mg OD	Consider 110mg BD	Reduce dose to 30mg OD
15-30	Use with caution		Contraindicated	
<15	Contraindicated	Contraindicated	Contraindicated	Contraindicated

Table 10. The use of DOACs in patients with reduced creatinine clearance.

Warfarin

- While it is becoming less common to use warfarin, it may still be used in patients who have a ongoing high risk of bleeding and may benefit from rapid reversal of anticoagulation. It may also be used to monitor patient's compliance, for patients with recurrent VTE whilst taking a DOAC, patients with a low bodyweight and for patients with a creatinine clearance of <15ml/min.
- Warfarin should always be used for patients who have had a PE and who are diagnosed with triple positive antiphospholipid syndrome.
- Warfarin may not be suitable for patients who are unable to comply with changes in dosage or regular INR checks.
- For pulmonary embolism, the standard target INR is 2-3.
- Induction treatment with warfarin should always follow a validated induction dosing algorithm; there are induction protocols available on the GGC Clinical Guideline platform ([link](#)). Dalteparin should be continued for at least 5 days after warfarin is started and the INR has been ≥ 2 for two consecutive days.
- Patients should be referred to the Anticoagulant Clinic at discharge ([link](#)).

Duration of Anticoagulation

Duration of anticoagulation should be assessed at a PE follow-up clinic; however, the below guidance may be used to inform patients of their likely duration at the time of discharge.

The length of ongoing anticoagulation should depend on the patient's VTE recurrence risk, taking into account the individual risk of bleeding. Clinical judgement should be used to assess an individual's bleeding risk, although the HAS-BLED and VTE-BLEED score may be useful.

Recommendations

The following are recommendations only and, in each case, clinical judgement and an individual's bleeding risk should be taken into consideration.

Scenario	Duration of Anticoagulation
All patients with PE	A minimum of 3 months
First episode of PE and active cancer	Indefinite or until cured
First episode of PE and single/dual positive Antiphospholipid syndrome	Indefinite (with apixaban)
First episode of PE and triple positive Antiphospholipid syndrome	Indefinite (with warfarin)
First episode of PE with no identifiable risk factor	Consider indefinite*
First episode of PE with a persistent risk factor	Consider indefinite*
First episode of PE with a minor transient risk factor	Consider indefinite*
First episode of PE secondary to a major transient/reversible risk factor	3 months
Recurrent VTE (more than one episode) not related to a transient/reversible risk factor	Indefinite*

*If using anticoagulation for > 3 months, and the patient doesn't have active cancer, continue anticoagulation with a DOAC at prophylactic dose (i.e. Apixaban 2.5mg BD, Rivaroxaban 10mg OD).

Risk of VTE Recurrence

Estimated risk for long-term recurrence	Risk factor category for index PE	Examples
Low (<3% per year)	Major transient or reversible factors associated with >10-fold increased risk for the index VTE event (compared to patients without the risk factor)	<ul style="list-style-type: none"> • Surgery with general anaesthesia for >30 min • Confined to bed in hospital (only "bathroom privileges") for ≥3 days due to an acute illness, or acute exacerbation of a chronic illness • Trauma with fractures
Intermediate (3-8% per year)	Transient or reversible factors associated with ≤10-fold increased risk for first (index) VTE	<ul style="list-style-type: none"> • Minor surgery (general anaesthesia for <30 min) • Admission to hospital for <3 days with an acute illness • Oestrogen therapy/contraception • Pregnancy or puerperium • Confined to bed out of hospital for ≥3 days with an acute illness • Leg injury (without fracture) associated with reduced mobility ≥3 days • Long-haul flight
	Non-malignant persistent risk factors	<ul style="list-style-type: none"> • Inflammatory bowel disease • Active autoimmune disease
	No identifiable risk factor	
High (>8% per year)		<ul style="list-style-type: none"> • Active cancer • One or more previous episodes of VTE in the absence of a major transient or reversible factor • Antiphospholipid antibody syndrome

Discharge and Follow Up

Discharge Check-list

- ☐ The patient is aware of the diagnosis of PE
- ☐ The patient is aware of the importance of continuing to take anti-coagulation for the treatment of acute PE and to prevent further PE
- ☐ The patient has been referred for follow-up (see below)
- ☐ The patient is aware that they will be seen in a follow-up clinic
- ☐ The patient has an adequate supply of anticoagulant
- ☐ The GP is informed of the PE and anticoagulation on the discharge letter

Follow Up

Follow-up should be performed for all patients with PE, regardless of severity and risk.

Follow-up should be performed at 3-6 months by a PE specialist. This is performed to help identify patients who are at risk of a post-PE syndrome, which includes pulmonary hypertension, and to help clarify duration of anticoagulation. Follow up includes an assessment of patient's symptoms and their risk factors for developing Chronic Thromboembolic Pulmonary Hypertension (CTEPH). Other investigations, such as an echocardiogram, NT-proBNP and V/Q scan may then be performed.

How to Follow Up

Currently, post-PE follow up will vary at each hospital across Greater Glasgow & Clyde. If no formal referral pathway exists, forward the discharge letter to the nominated respiratory consultant dealing with PE follow up and include pertinent information such as the risk status at diagnosis, the method of anticoagulation and whether thrombolysis was given.